

Physiological plasticity of *Arachis hypogaea* cultivars under drought conditions of semiarid region of India

KA KALARIYA^{1,2}, AL SINGH¹, RUPESH NAKAR^{1,3}, PV ZALA¹, CHAKRABORTY K^{1,4} and CB PATEL¹

¹ICAR Directorate of Groundnut Research, Junagadh, Gujarat 362 001, India; ²ICAR-Directorate of Medicinal and Aromatic Plants Research, Boriavi, Anand, Gujarat 387 310, India; ³Department of Botany, Sheth PT Arts and Science College, Godhra, Gujarat 389001, India; ⁴Central Rice Research Institute, Bidyadharpur, Cuttack, Odisha 753006; E-mail : kuldeep_ka@yahoo.co.in

(Received : July 13, 2017 ; Accepted : January 10, 2018)

ABSTRACT

Physiological plasticity of sixty peanut cultivars, belonging to four botanical groups, were evaluated and compared during *Kharif* season under well-watered (with protective irrigation; P) and natural drought (under rainfed; RF) conditions. Based on physiological and yield attributes to identify the promising genotypes. The days required for 50% flowering varied from 24.5-34.0 days and 26.0-37.7 days with an average of 28 and 30 days in P and RF crops, respectively. The natural drought (RF) condition delayed crop maturity (112-132 days) as against 113-119 maturity days in P. Interestingly, 30 cultivars matured within 113 days at 2130 °C degree days under both the condition indicating their adaptability and plasticity to drought. The mean pod yield of peanut cultivars were 1260 kg ha⁻¹ under P and 1130 kg ha⁻¹ under RF conditions, however cultivars ICGS 5, JGN 23, AK 265, GG 5, GG 11, GG 16, Girnar 1, AK 159, SBXI showed > 1300 kg ha⁻¹ pod yield under both the conditions. The cultivars with early flowering and maturity high SCMR, low SLA, high yield and HI showed the escape mechanism and were considered as most promising for rainfed. There is greater likelihood of drought situation. Our study showed, Spanish bunch (VUL) group was more suitable compared to Virginia bunch (HYP), Virginia runner (HIR) and Valencia (FST) peanut group for desirable traits in rainfed condition. The cultivars JGN 23, SB XI, and Girnar 1 showed most of the desirable characters with high physiological plasticity and hence, can be of immense use for rainfed conditions.

Key words: Degree days, Flower initiation, Natural drought, Peanut, Physiological Plasticity

The peanut is a major food legume of tropical and sub-tropical region of the globe and grown in about 110 countries under rainfed condition mostly with full of uncertainty of weather conditions (FAO, 2014, Singh *et al.* 2013). The productivity of peanut is less than 1000 kg ha⁻¹ in more than 30% of the peanut growing countries in the world, whereas it is between 1000-2000 kg ha⁻¹ in 40-45 % of the countries. Only 25% of the 110 countries possess productivity above 2000 kg ha⁻¹ (FAO, 2014). India though has the largest peanut area (5.53 m ha) in the world, but its average productivity is only around 1300 kg

ha⁻¹, which fluctuates between 990 to 1750 kg ha⁻¹ mainly due to its cultivation as rainfed crop without protective irrigation during kharif season (Singh 2004, 2011; Singh *et al.* 2013). However with good cultural practices and protective irrigations farmers are harvesting up to 4000 kg ha⁻¹ pod yield in certain areas (Singh, 2011). This calls for the attention of researchers to look into the matter and modify the recommendations.

Crop productivity per unit water is important especially for developing water use efficient cultivars (Codon *et al.* 2004). Physiological parameters associated with drought tolerance can be utilized for identification of drought resistant cultivars which can be used in crop improvement programmes (Nautiyal *et al.* 1999, 2012; Singh *et al.* 2013, 2014a, b). The leaf area and specific leaf area (Specific leaf area) are strongly correlated with photosynthesis and transpiration efficiency and thus dehydration tolerance capacity (Nageswara Rao *et al.* 1992; Wright *et al.* 1994). Chlorophyll is the major photosynthetic pigment and high chlorophyll density under water deficit stress is an indicator of tolerance (Arunyanark *et al.* 2008). The SLA and SPAD chlorophyll meter readings (SCMR) are quite useful, among the surrogate non-destructive traits as indirect selection tools for drought tolerance (Upadhyay, 2005; Nigam, 2008).

As majority of the peanut growing area belongs to semi-arid environment, information is required in making strategies for improvement of drought tolerant cultivars with high yield. In India, now there are more than 190 released peanut cultivars were released and out of these only 50-60 cultivars at the most in seed chain. There is hardly any consolidated report of testing all of these cultivars for drought tolerance under rainfed condition together. Plant response to drought are always dynamic in nature with respect to space, time, intensity of stress etc. Change in an organism's phenotype triggered by such variations is called phenotypic plasticity (Bradshaw, 1965). Many researchers have tried to assess the plasticity under challenging environments and found inverse relationship between tolerance and plasticity which was fit for fitness-related traits but was trait dependent

for underlying traits (Couso and Fernandez, 2012). We tried to measure the plasticity of these genotypes through analysing the performance of 60 peanut cultivars during kharif season under both protected and natural drought (rainfed) condition to find out the physiological parameters responsible for high yield and identification of cultivars with drought escape mechanism.

MATERIALS AND METHODS

Field experiment: A field experiment was conducted at the research farm of ICAR-Directorate of Groundnut Research, Junagadh, India, in a clayey calcareous vertisol soil having medium fertility in semi arid region of south Saurashtra. Sixty released peanut cultivars belonging to four different botanical groups (Table 1), comprising of 34 Spanish bunch (VUL), 16 Virginia bunch (HYP), 9 Virginia runners (HIR) and one Valencia (FST) were sown during Kharif season-2012 in a single row plot in screening blocks. The field was prepared and 40 kg N, 50 kg P and 50 kg K was applied in the soil as basal dose (Singh and Basu, 2005). All cultivars were sown in a 5 m row and having 45 x 10 cm spacing in three replications on 11th July, 2012 under sufficient moisture conditions. In the control plots (P) protected irrigation was provided whenever there was short fall in rain during the entire cropping season. On the other hand, in rainfed plot (RF) crop was raised under natural condition without any irrigation. The crop was harvested at physiological maturity of respective botanical groups.

Drought spell and its intensities during cropping season: In the present study, the cropping season spanned between 28th and 47th standard meteorological weeks in year 2012 during which the mean maximum and minimum temperature were 33.2 °C and 24.7°C, respectively, while

mean relative humidity was 72.1% and total bright sun shine hours was 594 (Table 2). The total rainfall from sowing till harvest was only 229 mm as against the total evaporation rate of 594 mm and there were, a total of three drought spells of various intensities observed during the entire cropping season in rain-fed crop. The crop faced the first drought spell of 10 days during 11 to 20 DAS with only 1.2% rainfall against the evaporative demand of 34.9 mm, the second drought spell of 20 days during 21 to 40 DAS with 13% rainfall against the evaporative demand of 58.5 mm and the last drought spell of 40 to 60 day starting from 72 DAS to harvest where the crop has received only 2.0 mm rainfall which was 0.6% of evaporative demand.

Flowering and morphological parameters: The days to flower initiation, 50% flowering, and total flowers produced during first 10 days were recorded in each cultivars under both the conditions. At 70 DAS three plants from each cultivars and treatments were uprooted and plant height, number of leaves on main axis, number of pods and pegs and dry biomass per plant was recorded. Keeping the base temperature uniformly at 10°C, the cumulative thermal time (CTT) expressed as θ (°C d, number of degree-days above the base temperature) required for initiation of flowering (θ_{If}), 50% flowering (θ_{f50}) and maturity (θ_m) were calculated by summation of daily mean temp minus 10 (Vasudeva et al. 1992).

SCMR, Leaf Area, Specific leaf area and yield parameters: The SCMR were recorded in the third fully expanded leaf facing sun from the top of the peanut plant at 70 DAS using SPAD-502 (Konika-Minolta, Japan) in each cultivar in triplicates. Twenty leaflets from third compound leaves were collected for measurement of leaf area, specific leaf area (SLA) and relative water content (RWC). Crop was harvested at maturity, dried in sun for a

Table 1. List of selected 60 peanut cultivars used in this study

S.N.	Cultivars	Habit group	Year of release	Area of adoption	Special characters
1	AK-159	VUL	2002	Maharashtra and Madhya pradesh (MP)	High oil content
2	Chico	VUL	--	--	Early maturity
3	DRG 12	VUL	1994	Andhra Pradesh(AP), Tamil Nadu(TN), Maharashtra, Karnataka	High yielding
4	DRG 1	VUL	--	--	--
5	GG 2	VUL	1983	Gujarat	Water use efficient
6	GG 20	HYB	1992	Gujarat	Large seeded with low aflatoxin contamination
7	GG 5	VUL	1999	Gujarat	Drought tolerant; leaflets stay green at maturity
8	GG 6	VUL	2003	Gujarat	Early maturity
9	GG 4	VUL	1993	Gujarat	High yielding and early maturity
10	GG 7	VUL	2001	Gujarat & southern Rajasthan	Early maturity
11	GG 8	VUL	2006	Northern Maharashtra and Madhya Pradesh	---
12	Girmar 1	VUL	1988	Western Maharashtra, T.N. and A.P.	Multiple diseases resistant, early maturity
13	Girmar 3	VUL	2010	West Bengal (WB), Orissa, Manipur	---
14	GPBD 4	VUL	2004	Maharashtra, Karnataka, AP & TN	---
15	ICGS 37	VUL	1990	Gujarat, northern Maharashtra and MP	Tolerant to end-of-season drought; photo-period insensitive

16	ICGS44	VUL	1988	Gujarat, northern Maharashtra & MP	High seed protein (25%) content
17	ICGV 86590	VUL	1991	Peninsular India	Multiple diseases resistant
18	ICGV 91114	VUL	2007	AP	Early maturity
19	JAL 42	VUL	--	--	Early maturity
20	JGN 23	VUL	2009	Madhya Pradesh	Drought tolerant
21	JL 24	VUL	1978	Maharashtra	Early maturity
22	JL 286	VUL	2004	Maharashtra	Early maturity
23	Kadiri5	VUL	2005	AP	Drought tolerant
24	Kadiri9	VUL	2009	AP	Tolerant to early and late season drought
25	PratapMungfali 1	VUL	2005	Rajasthan	Early
26	SB XI	VUL	1965	Maharashtra	Tolerant to <i>Aspergillusflavus</i> colonization
27	SG 99	VUL	2004	Punjab	---
28	TAG 24	VUL	1991	Maharashtra	Early maturity
29	TG 37A	VUL	2004	Gujarat, Rajasthan, Uttar Pradesh (UP), Orissa, W.B. Bihar and Assam	Possesses fresh seed dormancy (up to 15 days)
30	TG 51	VUL	2008	W.B., Orissa, Jharkhand and Assam	---
31	TLG 45	VUL	2004	Maharashtra	Large seeded
32	TMV 2	VUL	1940	Tamil Nadu, A.P. and Karnataka	Widely adapted
33	TPG-41	VUL	2004	All India	Large seeds, high O/L ratio, 25 day fresh seed dormancy
34	VRI 2	VUL	1989	Tamil Nadu	Tolerant to ELS, LLS & rust
35	CSMG 84-1	HYR	1992	Uttar Pradesh Rajasthan and Haryana	Variegated kernel colour
36	CSMG 9510	HYR	2005	U.P., Punjab, & northern Rajasthan	40-45 day fresh seed dormancy
37	DSG 1	HYR	1997	Karnataka	--
38	GG 11	HYR	1984	Gujarat	Resistant to rust
39	GG 16	HYR	2006	TN, AP, Kerala & southern Maharashtra	---
40	ICGV 88448*	HYR	--	--	Extra bold
41	M 13	HYR	1972	Punjab	Tolerant to leaf spots
42	M 335	HYR	1986	Punjab	Large seeded, Tolerant to ELS and LLS,
43	Somnath	HYR	1990	Gujarat and Rajasthan	Large seeded, early maturity,
44	AK 265	HYB	2007	Southern Maharashtra, Karnataka, AP, and TN	Drought tolerant
45	B 95	HYB	1993	Southern Maharashtra.	Large seeded, high yielding
46	BAU 13	HYB	1993	Bihar	Large seeded
47	CSMG 884	HYB	1999	UP, Punjab and Rajasthan	Large seeded, early maturity
48	DRG 17	HYB	1994	Rajasthan, Punjab, UP & Haryana	Tolerant to moisture stress
49	GG 14	HYB	2003	Northern Rajasthan, Punjab, Haryana & UP	
50	Girnar 2	HYB	2008	UP, Punjab, northern Rajasthan	Large seeded, stay green leaves at harvest
51	HNG 10	HYB	1998	Uttar Pradesh, Punjab, Rajasthan and Haryana	High yielding
52	HNG 69	HYB	2010	UP, Punjab and northern Rajasthan	High yielding
53	ICGS 5	HYB	1992	UP Rajasthan and Haryana	Drought tolerant
54	ICGS 76	HYB	1989	Southern Maharashtra and Karnataka	Resistant to ELS and LLS
55	ICGV 86031	HYB	--	--	Tolerant of iron chlorosis
56	ICGV 86325	HYB	1994	Southern Maharashtra, AP, Karnataka, Kerala, TN	High yielding
57	Kadiri 3	HYB	1978	AP	Clustered bearing
58	LGN 2	HYB	2001	Gujarat & southern Rajasthan	High yielding
59	TKG 19A	VUL	1995	Konkan region of Maharashtra	Bold and HPS grade kernels
60	Gangapuri	FST	1971	Madhya Pradesh	Moderately resistant to foliar disease

* the cultivars marked with are promising genotypes

Table 2. Weather parameters at various peanut crop growth stages at Junagadh, Gujarat India during the cropping season Kharif 2012

Growth Period	Crop growth stages	Temperature (°C)				RH(%)	BSS (h)	Evap (mm)	Rainfall (mm)	SMC at the end of the period
		Max	Min	mean						
1-10 DAS (11-20 July)	Cracking, establishment and early vegetative growth	33.1	25.9	29.5	76.4	5.8	34.9	24.3 (69.6)	9.6%	
11-20 DAS (21-30 July)	Vegetative growth	33.8	26.6	30.2	71.4	0.6	58.5	0.7 (1.2)	8.5%	
21-40 DAS (31 July to 19 Aug)	Vegetative growth, flower initiation and peg initiation	32.7	25.8	29.2	76.5	1.8	76.3	9.9 (13.0)	7.5%	
41 TO 67 DAS (20 Aug to 16 Sept)	Flowering, Peg initiation to beginning seed	31.1	24.7	27.9	85.5	21.1	62.2	262.4 (422)	7.5%	
68 TO 133 DAS (17 Sept to 20 Nov)	Beginning seed to full maturity and harvest	35.4	20.4	27.9	50.7	565.2	361.6	2.0 (0.6)	6.5%	
Mean		33.2	24.7	28.9	72.1					
Total						595	594	299 (50)		

Where, RH is Relative humidity, BSS is Bright sun shine hours during the period and the figures in parenthesis indicate percent rainfall of evaporative demand.

Table 3. Mean, minimum and maximum and standard deviation (SD) values of various parameters studies in 60 peanut cultivars

	Days for flowering Initiation		Days to 50% flowering		Total flowers produced in first 10 days		RWC at 70DAS		SCMR at 70DAS		SLA at 70DAS	
	P	RF	P	RF	P	RF	P	RF	P	RF	P	RF
Mean	23.2	26.3	27.7	31.2	61.7	57.3	95.6	84.9	37.5	39.5	185	147
Min	19.3	23	24.5	26	19	10	92.3	71.5	25.3	27.9	135	112
Max	28	32.5	34	37.7	111	105	97.1	91.8	44.4	50.4	293	210
SD	2.27	2.21	2.18	3	24.5	26	0.86	4.43	4.02	4.27	34	16

	Number of leaves on main axis		Plant height (cm)		Number of pods/plant		Pod yield kg ha ⁻¹		Fodder yield (kg ha ⁻¹)		HI	
	P	RF	P	RF	P	RF	P	RF	P	RF	P	RF
Mean	15.5	14.0	43.6	39.8	11.4	13.2	1260	1130	3330	2860	0.29	0.29
Min	13.0	11.0	30.8	28.1	7.9	7.4	530	390	930	1260	0.11	0.17
Max	18.7	17.2	56.4	55.3	18.4	18.1	2110	1700	6440	4520	0.41	0.42
SD	1.35	1.23	6.39	5.30	2.18	2.26	530	390	930	1260	0.11	0.17

Table 4. Analysis of variance among various physiological traits (Mean Sum Square values)

Effects	DF	Initiation of flowering (days)	50% flowering (days)	Total flowers	RWC (%)	SCMR	SLA
		Treatment	1	872.7**	1123.6**	1690**	10223.9**
Variety	59	26.2**	34.5**	3343.3**	29.9**	74.2**	3112**
Treatment x Variety	59	3.9**	6.8**	487.4**	31.4**	28.9**	1221**
Residual	240	2.3	3.6	0.6	3	9.6	78
Total	359						

Effects	DF	No. of leaves on main axis	Plant Height	No. of pods/plant	Pod wt/plant	Foddrwt/plant	HI
Treatment	1	210.8**	1336.9**	285.2	32.3*	162**	5.53391
Variety	59	7.6**	193.3**	21.9	21.8**	197.3**	0.01652**
Treatment x Variety	59	2.4**	13.5	7.7	5.7	23.1	0.00283*
Residual	240	1.5	12.2	7.1	5.5	19.3	0.00202
Total	359						

* and ** indicates significance at 0.05 and 0.01 level (P value < 0.05, 0.01)

Table 5. Peanut cultivars with desirable physiological traits under rainfed and protected conditions during Kharif2012

Traits	Protected	Rainfed
	Desirable cultivars	Desirable cultivars
Pod yield	ICGS 5, JGN 23, AK 265, GG 5, GG 11, GG 16, GG 20, Girnar 1, AK 159, ICGV 86325, CSMG 9510, HNG 10, M 13, BAU 13, JAL 42, SB XI, DSG 1 (>1450 kg ha ⁻¹)	ICGS 5, JGN 23, AK 265, GG 5, GG 6, GG 7, GG 11, GG 16, Girnar 1, Gangapuri, AK 159, SBXI, TMV 2, DRG 1, DRG 12, JL 286 (>1300 kg ha ⁻¹)
HI	TG 51, JAL 42, JGN 23, Girnar 1, SB XI, ICGS 44, GG 2, TAG 24, JL 286, JL 24, TG 37A, ICGS 5, DRG 1, HNG 10, (> 0.33)	TG 51, JAL 42, JGN 23, Girnar 1, SB XI, ICGS 44, GG 2, TAG 24, JL 286, GG 5, GG 7, DRG 12, TLG 45, ICGS 37, DRG 1, Gangapuri, TMV 2 (> 0.33)
SCMR	GG 20, DRG 17, ICGV 86031, ICGS 37, ICGS 44, CSMG 9510, GG8, B 95, SG 99, Somnath, HNG 69, Kadiri 9, TMV 2, TLG 45, CSMG 884, BAU 13, Kadiri3 (>40)	TPG 41, SG 99, DRG 12, Kadiri 9, GG 7, ICGV 86325, LGN 2, ICGS 44, ICGV 86031, GG 11, DRG 17, ICGV 86590, JL 24, ICGS 5, ICGS 37, M 335, TG 37A, Kadiri 5, TKG 19A, GG 8, B 95, Somnath, CSMG 9510, GG 20 (>40)
SLA	ICGS 37, ICGV 86031, ICGV 86590, ICGV 88448, CSMG 884, Somnath, GG 11, M 13, ICGS 44, B 95, TPG41, ICGV 91114 (<160 cm ² g ⁻¹)	ICGS 37, ICGV 86031, ICGV 86590, ICGV 88448, CSMG 884, Somnath, GG 11, M 13, ICGS 44, TKG 19A, Girnar 2, Kadiri 9, DRG 12, CSMG 9510, DSG 1, TAG 24, DRG 17, SG 99, Girnar 3, HNG 10, M 335, GG 14, ICGV 86325, Pratapmungfali 1 (<141 cm ² g ⁻¹)
Early Flowering	JGN 23, GG 2, GG 4, GG 7, GG 8, Girnar 1, JL 24, JL 286, Kadiri 5, JAL 42, SB XI, AK 159, Gangapuri, TMV 2, Chico, DRG 1, TAG 24, ICGS 37, ICGV 91114, TKG 19A, (within 26 days at 506 C°D)	JGN 23, GG 2, GG 6, GG 7, GG 8, Girnar 1, JL 24, JL 286, Kadiri 5, JAL 42, SB XI, AK 159, Gangapuri, TMV 2, Chico, Pratapmungfali 1, GPBD 4, ICGS 5, TG 51, TLG 45, SG 99 (within 30 days at 565 C°D)
Early maturity	Chico, JGN23, GG 2, GG 4, GG 5, GG 6, GG7, GG8, Girnar 1, JL 24, JL 286, Gangapuri, TMV2, SB XI, AK 159, Pratapmungfali1, ICGS 5, ICGS 37, ICGS 44, ICGV 91114, ICGV 86031, JAL 42, DRG 1, TAG 24, TG 37A, TPG41, TLG 45, TG51, VRI 2, Kadiri5, (within 113 days at 2130 C°D)	Chico, JGN23, GG 2, GG 4, GG 5, GG 6, GG7, GG8, Girnar 1, JL 24, JL 286, Gangapuri, TMV2, SB XI, AK 159, Pratapmungfali1, ICGS 5, ICGS 37, ICGS 44, ICGV 91114, ICGV 86031, JAL 42, DRG 1, TAG 24, TG 37A, TPG41, TLG 45, TG51, VRI 2, Kadiri5, GPBD 4, SG 99, (within 112 days at 2114 C°D)

Table 6. Yield trait wise classification of cultivars for drought tolerance

Trait and trait combinations	Treatment conditions	List of cultivars
High yield and HI and early flowering,	P	JGN23, Girnar 1, JAL 42, SB XI
	RF	JGN 23, GG 7, Girnar 1, SB XI, JL 286, GG 11, Gangapuri, TMV 2
High yield and SCMR	P	GG 20, CSMG 9510, BAU13
	RF	ICGS 5
High SCMR and Low SLA	P	ICGV 86031, ICGS 37, ICGS 44, CSMG 884 B 95, Somnath
	RF	Kadiri 9, SG 99, ICGV 86031, ICGV 86325, ICGV 86590, Somnath, GG 11, M 335, TKG 19A, ICGS 37, CSMG 9510
High yield and early Maturity	P	ICGS 5, JGN 23, GG 5, AK 159, JAL 42, SB XI, Girnar 1,
	RF	ICGS 5, JGN 23, GG 5, GG 6, GG 7, Girnar 1, Gangapuri, AK 159, SBXI, TMV 2, DRG 1, JL 286
High yield and HI, and early flowering and maturity	P	JGN23, Girnar 1, JAL 42, SB XI
	RF	JGN23, Girnar 1, SB XI

P and RF are protected and Natural drought (rainfed) treatments, respectively.

week and pod and haulm yields, HI and other post-harvest observation was recorded.

Statistical analysis: All the data were subjected to statistical analysis following Gomez and Gomez (1984). Linear correlation was worked out between various physiological parameters studied under protected and rain-fed conditions. One-way ANOVA was also carried out for both the treatments using DSTAAT software.

RESULTS AND DISCUSSION

The drought caused initial plant death, delayed

flowering and reduced leaf area, SLA, plant height, number of flowers and harvest index, but increased SCMR under rain-fed condition. Significant variation among cultivars was observed for various physiological parameters the details of which are discussed below.

Flowering: In peanut cultivars, the initiation of flowering started from 19 to 28 DAS under protected condition with an average at 23 DAS, but under RF the average flower initiation time was 26 DAS i.e. delayed by 3 days (Table 3). Accordingly, the 50% flowering time was attained in 28 DAS under P and 31 DAS in RF conditions which corresponded to cumulative thermal time θ_f of 546

and 602 °C d, respectively. Thus mean 50% flowering was also delayed by 3-4 days in RF condition. We found, 20 cultivars flowered within 26 days at θ_f of 506 °C d under P conditions, whereas 21 cultivars flowered within 30 days at θ_f of 565 °C d in RF condition. Interestingly 14 cultivars were common under both the situations. The numbers of flowers produced from the day of 50% flowering to next 10 days showed very high variation among cultivars. It ranged from 19 (TPG 41) to 111 (HNG 69) flowers per plant under P and 10 (JAL 286) to 105 (BAU 13) under RF.

RWC, SCMR and SLA: The RWC, SCMR and SLA measured at 70 DAS showed significant variation within the cultivar as well as under different treatment conditions (Table 3). The mean RWC value of these cultivars was 95.6 (with a range of 92.3 to 97.1) under P conditions which decreased to 84.9 (with a range of 71.5 to 91.8) under RF condition. Interestingly, there were 35 cultivars showing >85 RWC under RF conditions. The mean SCMR value of these cultivars was 37.5 under P conditions which increased to 39.5 under RF. Seventeen cultivars under P and 24 cultivars in RF showed higher SCMR values of >40 and of these 10 cultivars were common under both the conditions (Table 5). The mean SLA was 185 and 147 cm²g⁻¹ under P and RF conditions, respectively. Among the cultivars highest SLA was observed in VRI 2 (293 cm²g⁻¹) and lowest in ICGS 37 (135 cm²g⁻¹) in P however under RF the SLA was highest in Chico (210 cm²g⁻¹) and lowest in Kadiri 9 (112 cm²g⁻¹). Twelve cultivars under P showed SLA values less than 160 cm²g⁻¹, while 24 cultivars showed SLA values less than 140 cm²g⁻¹ under RF condition indicating drought adaptive response in these cultivars (Table 5).

Plant height, number of leaves and pods: The morphological characters varied significantly with the treatment conditions (Table 4). Under protected condition mean plant height was 43.6 cm, which decreased to 39.8 cm under rainfed condition. Similarly, the leaves on main axis was reduced from 16 to 14 under RF conditions (Table 3). Under protected condition the number of cultivars with more number of leaves on main axis was higher but, under RF there were equal no. of cultivars showing less and more leaves. On an average the cultivars GG 2, JGN 23, LNG 2, GG 7, GG 14, DRG 12 and TG 51 showed more leaves.

On an average there was 11.6 pods plant⁻¹ in P and 14 in RF at 70 DAS and out of 60 cultivars, nine showed more number of pods under P condition and only seven under RF condition. The cultivars TPG 41, Gangapuri, DRG 14 and TLG 45 produced more number of pods. Five cultivars which showed more pods under P did not maintain same trend under RF. Interestingly, three cultivars SB IX, JGN 23 and Pratap Mungfali 1 showed better response with more number of pods under RF.

Maturity period and observations at harvest: There was a considerable effect of drought on the overall maturity of the crop as a result the cultivars under RF condition took 112-132 days with a CTT of 2114-2420 °C d to mature as against 113-119 days with 2130-2224 °C d in protected condition. Interestingly 33 cultivars matured within 112 days at 2114 °C d under RF condition and 30 cultivars matured within 113 days at 2130 °C d under P condition and more so 30 cultivars were common in both the condition indicating their adaptability and plasticity to drought.

There were significant differences in the mean pod yield of peanut under protected (1260 kg ha⁻¹) and RF (1130 kg ha⁻¹) conditions. Seventeen cultivars showed > 1450 kg ha⁻¹ pod yield under P, however 16 cultivars showed > 1300 kg ha⁻¹ pod yield under RF condition and of these nine were common in both the conditions (Table 5). Due to continuous drought there was death in plant under RF conditions, but no death under protected condition. As a result the average number of pods plant⁻¹ increased under RF condition due to thinning of plant population and it was 11.4 under P and 13.2 pods plant⁻¹ in RF condition. There was drastic reduction in the haulm yield of peanut under RF (2860 kg ha⁻¹) as compared to the one under protected (3330 kg ha⁻¹) conditions. Fifteen cultivars showed >4000 kg ha⁻¹ haulm yield under P, however under RF condition only 12 cultivars could produce > 3300 kg ha⁻¹ haulm yield.

In the present agriculture, for efficient use of water our focus should be to raise water use efficient cultivars both under irrigated as well as rain-fed conditions (Codon *et al.* 2004). Though traits conferring capacity of dehydration avoidance and tolerance are available, integrated traits expressing tolerance at organ level are more useful (Singh 2011, Singh *et al.* 2013). Selection based on genetics, yield and physiology is part of physiological genetic approach (Reddy *et al.* 2003, Singh *et al.* 2010). Flowering in peanut starts at 20 DAS with effective flowering observed at 30 DAS (Singh 2011). But, there is diversity in maturing pods due to extended flowering period and pod yield depends upon flower production (Singh, 2004, 2011). In this study, significant variation among cultivars was found for most of the studied traits. Under rainfed condition, there was delay in flowering due to insufficient rainfall during 11-19 DAS however; during 20-31 DAS there was scanty rainfall (~10 mm), which initiated flowering with production of more number of flowers plant⁻¹ day⁻¹ on an average. In present study, though 50% flowering was observed at 25-34 DAS under protected (P) and 26-38 DAS under rain-fed (RF) condition, but the effective flowering was observed between 28-38 DAS under P and 31-40 DAS under RF. We identified this as the critical yield determining stage which should not face drought.

Screening based on various physiological and agronomical traits resulted in identification of cultivars possessing desirable traits. In general the early flowering, high SCMR, pod yield and HI, and low SLA were identified as the desirable traits under both the conditions. The cultivars showing early flowering, high pod yield and high HI, high yield and SCMR, high SCMR and low SLA and high pod yield and early maturity in both the treatments are listed in Table 6. Further, the cultivars with early flowering, high SCMR, low SLA and high yield and HI showed the escape mechanism and were considered as highly suitable for rain-fed cultivation under drought situation however, the one showing *vice versa* physiological traits were most susceptible. Though a number of cultivars having desirable traits under both protected and rain-fed conditions were listed in Table 5, the cultivars JGN 23, SB XI and Girnar 1 showed most of the desirable characters and hence can be of immense use as donor parents for rain-fed conditions.

Reduced SLA provides lesser surface area for harvesting photosynthetic light, a protective mechanism of the photosynthetic pigments under stress condition as plants are not able to utilize all the absorbed photons and the unutilized photons is diverted towards the non photochemical quenching through heat generation. Also the reduced SLA provide a lesser leaf surface area in direct contact with the ambient air circulation which causes a loss of water from leaves. Wunna *et al.* (2009) reported positive association between SLA and HI in drought condition.

In this study, we found SCMR as a highly useful trait in identification of cultivars for drought tolerance. Bootang *et al.* (2010) reported that physiological parameters SCMR and SLW gave higher contribution to biomass under drought than pod yield and the HI and number of mature pods contributed to high pod yield. However, in this study, we found SLA and HI were highly useful traits for identifying cultivars under natural drought events.

Here in this study, the rain-fed crop faced three distinctly different drought spells from 11-20 DAS, 21-40 DAS and 68 DAS to harvest which resulted in delayed flowering, and lesser flowers production in first 10 days and affected yield and yield attributes. Seventeen cultivars showed > 1450 kg ha⁻¹ pod yield under P, however 16 cultivars showed > 1300 kg ha⁻¹ pod yield under RF condition and of these nine cultivars were common in both the conditions indicating their plasticity to drought stress. Earlier Nautiyal, *et al.* (2002) reported early stage drought in peanut does not affect yield, biomass production and nodule dry weight. On the contrary, end season drought, in various peanut cultivars, increased SCMR and SLW, but reduced biomass, pod yield and seed size without affecting the HI and number of (Bootang *et al.* 2010). The mean HI in this study under both facing

and RF treatment was similar. However, imposition of drought under rain-fed condition increased SCMR, but decreased RWC, number of flowers and pods and yield. Recently, SCMR has been found a more pertinent trait than SLA in summer peanut under transient water deficit stress condition (Kalariya *et al.* 2015a).

Among the four botanical groups, Spanish bunch group was found best with desirable traits in rainfed drought-prone condition. The cultivars like ICGS 5, JGN 23, AK 265, GG 5, GG 11, GG 16, Girnar 1, AK 159, SB XI showed > 1300 kg ha⁻¹ pod yield under both the conditions and found suitable for rain-fed cultivation. Combination of high SCMR with low SLA, high HI and low SLA, high yield and HI and early flowering behaviour are required high physiological plasticity for drought tolerance and hence are ideal for both protected and rain-fed cultivation. Cultivar screened for natural drought is of immense use in the areas where drought is occurs more often than not in semi-arid regions of the world. The study conclude that the cultivars with early flowering, high SCMR, low SLA, high pod yield and HI along with early maturity trait possess drought tolerance mechanism and were considered as highly suitable for rain-fed cultivation. Among different habit groups, Spanish bunch group was found more suitable. The cultivars *viz.* JGN 23, SB XI, and Girnar 1 showed most of the desirable characters indicating high physiological plasticity and hence can be of immense use for rain-fed cultivation.

REFERENCES

- Arunyanark A, Jogloy S, Akkasaeng C, Vorasoot N, Kesmla T, Nageswara Rao RC, Wright GC and Patanathai A. 2008. Chlorophyll stability is an indicator of drought tolerance in peanut. - Journal of Agronomy and Crop Science. **194**: 113-125
- Babitha M, Sudhakar P, Latha P, Reddy PV and Vasanthi RP. 2010. Screening of groundnut genotypes for water use efficiency and temperature tolerance. Indian Journal of Plant Physiology **11(1)**: 63-74.
- Bradshaw AD. 1965. Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics **13**: 115-155
- Bootang S, Songasri P, Jogloy S, Akkasaeng C, Vorasoot N, Tantisuwichwong and Potanathi A, 2010. Evaluation of peanut cultivars commonly grown in Thailand under water limited conditions. Asian Journal of Plant Science **9(6)**: 320-328
- Couso LL and Fernandez RJ. 2012. Phenotypic plasticity as an index of drought tolerance in three Patagonian steppe grasses. Annals of Botany **110**: 849-857.
- Codon AG, Richards RA, Rebetzke GJ and Farquhar GD. 2004. Breeding for high water use efficiency. Journal of Experimental Botany. **55**: 2447-2460
- FAOSTAT. 2014. Statistics Division
- Girdthai T, Jogloy S, Vorasoot N, Akkasaeng C, Wongkew, S, Holbrook, CC and Potanathi A, 2010. Association between Associations between physiological traits for drought tolerance and aflatoxin contamination in peanut genotypes under terminal drought. - Plant Breeding **129**: 693-699
- Gomez KA and Gomez AA. 1984. (Ed). Statistical procedure in Agriculture Research. -Wiley Publications, New York. pp: 680

- Hemidou F, Halilou O and Vandez V.2012. Assessment of groundnut under combined heat and drought stress. *Journal of Agronomy and Crop science* **10**: 1437-1439
- Kalariya KA, Singh AL, Chakraborty K, Zala PV and Patel CB. 2013. Photosynthetic characteristics of groundnut (*Arachis hypogaea* L.) under water deficit stress. *Indian Journal of Plant Physiology* **18**(2): 157-163
- Kalariya KA, Singh AL, Chakraborty K, Ajay BC, Zala PV, Patel CB, Nakar RN, Nisha Goshwami and Deepti Mehta. 2015a. SCMR: a more pertinent trait than sla in peanut genotypes under transient water deficit stress during summer. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* DOI 10.1007/s40011-015-0636-4
- Kalariya KA, Singh AL, Nisha Goshwami, Deepti Mehta, Mahatma M.K, Ajay BC, Chakraborty K, Zala PV, Vidhya Chaudhary, Patel Photosynthetic characteristics of peanut genotypes under excessand deficit irrigation during summer. *Physiology and Molecular Biology of Plants* DOI 10.1007/s12298-015-0300-8
- Nageswara Rao, RC Reddy LJ, Mehan VK, Nigam SN and McDonald D. 1992. Drought research on groundnut at ICRISAT. In: *Groundnut- A global Perspective: Proceeding of an International Workshop*, 25-29 Nov 1991, ICRISAT Asia Centre, Patancheru, Andhra Pradesh., India
- Nigam SN and Aruna R. 2008. Stability of soil plant analytical development (SPAD) chlorophyll meter reading (SCMR) and specific leaf area (SLA) and their association across varying soil moisture stress conditions in groundnut (*Arachis hypogaea* L.). *Euphytica*. **160**: 111-117
- Nautiyal PC, Ravindra V, Zala PV and Joshi YC. 1998. Enhancement of yield in groundnut following the imposition of transient soil-moisture-deficit stress during the vegetative phase. *Experimental Agriculture* **35**: 371-385
- Nautiyal PC, Ravindra V, Rathna kumar AL, Ajay BC and Zala PV. 2012. Genetic variations in photosynthetic rate, podyield and yield components in Spanish groundnut cultivars during three cropping seasons. *Field Crops Research* **125**: 83-91
- Reddy TY, Reddy VR and Anbumozhi V. 2003. Physiological responses of groundnut (*Arachis hypogaea* L.) to drought stress and its amelioration: a critical review. *Plant Growth Regulation* **41**: 75-88
- Rowland DL, Beasley JP Jr and Faircloth WH. 2010. Genotypic Differences in Current Peanut (*Arachis hypogaea* L.) Cultivars in phenology and Stability of these traits under different irrigation scheduling methods. *Peanut Science*. **37**: 110-123
- Saha RR, Aziz A, Begum F, Ahmed IM and Golder PC. 2010. Study on flowering and pod development pattern in seed production of groundnut. *SAARC Journal of Agriculture*. **8**(2): 11-18
- Samdur MY, Singh AL, Mathur RK, Manivel P, Chikani BM, Gor HK and Khan MA. 2000. Field evaluation of chlorophyll meter for screening groundnut (*Arachis hypogaea* L.) genotype tolerant to iron deficiency chlorosis. *Current Science* **79**: 211-214
- Sheshshayee MS, Bindu Madhava M, Rachaputi NR, Prasad TG, Udaykumar M, Wright GC and Nigam SN. 2006. Leaf Chlorophyll concentration relates to transpiration efficiency in peanut. *Annals of Applied Biology* **148**: 7-15
- Singh AL, Nakar RN, Goswami N, Mehta D, Subhangi Oza, Kalariya KA, Chakraborty K, Vidhya Chaudhari and Patel CB. 2013b. FYM and fertilizer increases photosynthetic efficiency and fluorescence in groundnut. In *Current Trends in Plant Biology Research* pp. 571-572
- Singh AL. 2004. Growth and physiology of groundnut. In M. S. Basu, N.B. Singh (Eds.): *Groundnut Research in India*. pp. 178-212. Junagadh, National Res Centre for Groundnut, ICAR
- Singh AL. 2011. Physiological basis for realizing yield potentials in groundnut. In A. Hemantranjan (Ed.): *Advances in Plant Physiology* Vol. 12. pp. 131-242 Scientific Publishers, Jodhpur- India
- Singh AL and Basu MS. 2005. Integrated nutrient management in groundnut-a farmer's manual. National Research Center for groundnut, Junagadh, India. 54 p
- Singh AL and Joshi YC. 1993. Comparative studies on the chlorophyll content, growth, N uptake and yield of groundnut varieties of different habit groups. *Oleagineux* **48**: 27-34
- Singh AL, Nakar RN, Goswami Nisha, Kalariya KA, Chakraborty K and Singh M. 2013. Water deficit stress and its management in groundnut (*Arachis hypogaea* L.) In A. Hemantranjan (Ed.): *Advances in Plant Physiology*. Vol. 14, pp. 375-465. Scientific Publishers, Jodhpur-India
- Singh AL, Nisha Goswami, Nakar RN, Kalariya KA and Chakraborty K. 2014a. Physiology of groundnut under water deficit stress. In A.L. Singh (Ed) *Recent Advances in Crop Physiology*, Vol. 1 pp.1-85. Astral International, New Delhi, India
- Singh AL, Nakar RN, Chakraborty K and Kalariya KA. 2014b. Physiological efficiencies of minicore peanut germplasm accessions. *Photosynthetica* **52**(4): 627-634
- Singh AL, Nautiyal PC and Zala PV. 1998. Growth and yield of groundnut (*Arachis hypogaea* L.) varieties as influenced by seed size. *Tropical Science* **38**: 48-56
- Singh S, Singh AL, Kalpana S and Misra S. 2010. Genetic diversity for growth, Yield and Quality traits in groundnut (*Arachis hypogaea* L.). *Indian Journal of Plant Physiology* **15**: 267-271
- Upadhyaya HD. 2005. Variability for drought resistance related traits in the mini core collection of peanut *Crop Science* **45**: 1432-1440
- Upadhyay HD, Sharma S, Singh S and Singh M. 2011. Inheritance of drought resistance related traits in two crosses in groundnut. (*Arachis hypogaea* L.). *Euphytica* **177**: 55-66
- Vasudeva MJ, Nigam SN and Huda AKS. 1992. The thermal time concept as a selection criterion for earliness in peanut. *Peanut Science* **19**: 7-10
- Wright GC, Nageswara Rao RC and Farquhar GD. 1994. Water use efficiency and carbon isotope discrimination in pea nut under water deficit conditions. *Crop Science*. **34**: 92-97